



Cholesterol Lowering Effect of *Cnidoscoulous aconitifolius* Leave Extracts in Egg Yolk Induced Hypercholesterolemia in Rabbit

Mathew Folaranmi Olaniyan^{1*}, Donatus F. Ozuaruoke² and Temitayo Afolabi¹

¹*Department of Medical Laboratory Science, Achievers University, Owo, Nigeria.*

²*Department of Education, Medical Laboratory Science Council of Nigeria, Abuja, Nigeria.*

Authors' contributions

This work was carried out in collaboration between all authors. Author MFO designed the study. Author DFO performed the statistical analysis. Author MFO wrote the protocol and wrote the first draft of the manuscript. Authors TA and MFO managed the analyses of the study. Authors DFO and TA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2017/35431

Editor(s):

(1) Thomas I. Nathaniel, University of South Carolina, School of Medicine-Greenville, Greenville, USA.

Reviewers:

(1) Eduardo Padilla Camberos, CIATEJ, Mexico.

(2) Siham Mostafa Ali El-Shenawy, National Research Center (NRC), Egypt.

Complete Peer review History: <http://www.sciencedomain.org/review-history/20197>

Original Research Article

Received 12th July 2017
Accepted 19th July 2017
Published 25th July 2017

ABSTRACT

Study Background: *Cnidoscoulous aconitifolius* leaf (Chaya; Efo Iyana ipaja in Yoruba language) is a common vegetable plant served as sauce and has many traditional health benefit claims such as treatment of alcoholism, insomnia, gout, scorpion stings, memory and vision impairment owing to its phytochemical and phytonutrient constituents.

Aim and Objectives: This work was designed to determine the cholesterol lowering effect of *Cnidoscoulous aconitifolius* in rabbits induced with hypercholesterolemia using egg yolk.

Materials and Methods: Fifteen (15) rabbits grouped into 3 major classes of 5 rabbits each has A (Control rabbits given only normal meal and water), B1 (5 rabbits give 20% egg yolk of normal meal and water for 7 days), B2 (B1 given 400 mg/kgBW of ethanolic extract of *Cnidoscoulous aconitifolius* leaf after egg yolk containing meal), C1 (5 rabbits give 20% egg yolk of normal meal and water for 7 days), C2 (C1 given 400 mg/kgBW of aqueous extract of *Cnidoscoulous aconitifolius* leaf after egg yolk containing meal). Plasma Total cholesterol(CHOL-T), Low Density Lipoprotein-

*Corresponding author: E-mail: olaniyanmat@yahoo.com;

cholesterol(LDL-C),Total Triglycerides(TG-T), High Density Lipoprotein-cholesterol (HDL-C) were evaluated in the rabbits by autoanalysis using ROCHE reagent on COBAS C111 auto Chemistry analyzer.

Results: There was a significant increase in the plasma CHOL-T, LDL-C, TG-T, and HDL-C in the rabbits given 20% of powdered egg yolk of the total meal weight and water for seven days compared with the control rabbits ($p<0.05$). There was also a significant decrease in the plasma value of CHOL-T, LDL-C and TG-T when the hypercholesterolemic rabbits (B1 and C1) were supplemented with 400 mg/kgBW of ethanolic and aqueous of extract *Cnidoscoulous aconitifolius* leaf for 7 days (B2 and C2) ($p<0.05$).

Conclusion: Induction of hypercholesterolemia using 20% egg yolk of normal meal and water resulted into a significant increase in plasma value of CHOL-T, LDL-C, TG-T, and HDL-C while supplements of 400 mg/kgBW of ethanolic and aqueous extract of *Cnidoscoulous aconitifolius* leaf for 7 days significantly reduced the plasma value of CHOL-T, LDL-C and TG-T in the hypercholesterolemic rabbits.

Keywords: Cholesterol; *Cnidoscoulous aconitifolius*; rabbits; hypercholesterolemia; egg yolk.

1. INTRODUCTION

Cnidoscoulous aconitifolius leaf (Chaya; Efo Iyana ipaja in Yoruba language) is a common vegetable South-Western part of Nigeria prepared as a sauce. It has many non-scientific traditional uses to cure alcoholism, insomnia, gout, scorpion stings, memory and vision impairment. *Cnidoscoulous aconitifolius* leaf is a good source of protein, vitamins, calcium, and iron; and is also a rich source of antioxidants [1]. However, raw leaves of the vegetable are toxic as they contain a glucoside that can release toxic cyanide (toxic hydrocyanic acid) similar to cassava. Cooking for about 5 minutes or more prior to consumption could inactivate the toxic components [1-3]. The leaf contains: protein (5.7%), crude fiber (1.9%), calcium (199.4 mg/100 g), potassium (217.2 mg/100 g), iron (11.4 mg/100 g), vitamin C (164.7 mg/100 g), and carotene (0.085 mg/100 g) [4]. Omotoso et al. [5] reported 20 phyto-chemotypes with different therapeutic activities, which include, 9-Octadecenoic acid (Z) and its esters, n-Hexadecanoic acid, n-Octadecanoic acid, n-Octacosane, 1,2,3-Propanetriol and its derivatives, and l-(+)-Ascorbic acid-2,6-dihexadecanoate. The Preliminary phytochemical analysis of *Cnidoscoulous aconitifolius* leaf reported by Obichi et al. [6] showed that the leaf extracts contain tannins (5.72 ± 0.00), saponins (12.49 ± 0.021), alkaloids (17.45 ± 0.65), flavonoids (23.72 ± 0.02), cyanogenic glycosides (0.75 ± 0.10) and phytate (1.97 ± 0.06) and concentration of vitamin A (5.24 mg/kg), vitamin B3 (1.40 mg/kg), vitamin B6 (37.23 mg/kg), vitamin B12 (15.98 mg/kg), vitamin C (382.00 mg/kg) and vitamin E (18.28 mg/kg).

A large egg of 50 g total weight, contains 17 g of egg yolk which contains about 2.7 g protein, 210 mg cholesterol, 0.61 g carbohydrates, and 4.51 g total fat. It contains fat-soluble vitamins (A, D, E, and K) are found in the egg yolk. Egg yolk is rich cholesterol and fat soluble vitamins. Rabbits are known as an animal good at storing fats or lipids [7-10].

This work was designed to determine the cholesterol lowering effect of *Cnidoscoulous aconitifolius* in rabbits induced with hypercholesterolemia using egg yolk.

2. MATERIALS AND METHODS

2.1 Study Area

Animal house of Achievers University, Owo, Nigeria equidistant between Nigeria Federal capital territory-Abuja and former Federal capital-Lagos. It has Latitude: 6.98575, Longitude: 5.27103 and Time Zone: UTC+1, Africa/Lagos.

Study population Rabbits were bought from Ojalkokoamajor Owo market and were identified and confirmed having same sex in the Department of Biological Sciences, Achievers University, Owo-Nigeria. These include 25 rabbits of the same sex with weight ranging from 1.0-1.4 Kg grouped as follows:

Group A: Five rabbits weighing 1.2 ± 0.1 Kg fed with normal meal and water for 7 days were studied as control group A.

Group B₁: Five rabbits weighing 1.1 ± 0.1 Kg fed with normal meal containing 20% of

powdered egg yolk of the total meal weight and water for seven days.

Group B₂: Five rabbits weighing 1.3 ± 0.1 Kg fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days followed by the administration of with 400 mg/Kg of ethanolic extract of *Cnidoscopus aconitifolius* for another seven days.

Group C₁: Five rabbits weighing 1.2 ± 0.1 Kg fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days.

Group C₂: Five rabbits weighing 1.2 ± 0.1 Kg fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days followed by the administration of 400 mg/Kg of aqueous extract of *Cnidoscopus aconitifolius* for another seven days.

2.2 Preparation of the *Cnidoscopus aconitifolius* Extracts

Cnidoscopus aconitifolius (iyana ipaja In Yoruba) was purchased from Ojalkoko, Owo-Nigeria and was identified by the Department of Biological Sciences. The leaves of *Cnidoscopus aconitifolius* (iyana ipaja In Yoruba)- *Cnidoscopus aconitifolius* was air dried for 14 days, Ethanolic and aqueous extraction was carried out by soaking 50 g of powers of *Cnidoscopus aconitifolius* into 500 ml of each of ethanol and sterile distilled water for 24 hours. Following the report of Das et al. [11] that solvent to sample ratio of 10:1 (v/w; solvent to dry weight ratio) has been used as ideal. Each extract was filtered through Whatmann filter paper No.1 and filtrates concentrated at room temperature in order to reduce the volume. Further concentration and drying by volume extraction was carried out using rotary evaporator and stored in refrigerator prior to use. Four hundred milligramme of the extract powder was dissolved in 2 ml of distilled water for administration.

2.3 Preparation of Egg Yolk Powder

Local eggs were purchased from Ojalkoko, Owo-Nigeria and presented to the Biological sciences department of Achievers University, Owo, Nigeria for identification. The shell of the egg was removed and the egg yolk was extracted. The egg yolk was air dried and grinded into powder.

2.4 Preparation of 20% Egg Yolk Powder of Normal Rabbit Meal

The normal meal was weighed. 20% of the weight was removed using weighing balance the 20% was replaced by egg yolk powder and this was used to induce hypercholesterolemia observed in B1 and C1.

2.5 Blood Specimen

Blood samples were collected from the veins lining the ear of the rabbits after each treatment into lithium heparinized bottles for the estimation of Total cholesterol, LDL cholesterol and Total triglycerides.

2.6 Determination of Biochemical Parameters

Plasma concentration of Total cholesterol, Low Density Lipoprotein-cholesterol (LDLCL), Total Triglycerides, High Density Lipoproteincholesterol (HDLCL) was determined by CABAS C111 auto-Chemistry analyzer using Roche reagent.

2.7 Ethical Consideration

Ethical guidelines on the use of rabbits for research work were obtained from the Department of Biological Sciences, Achievers University, Owo-Nigeria which was strictly adhered to.

2.8 Method of Data Analysis

The results obtained was subjected to statistical analysis using SPSS 18.0 to determine student "t" test, probability and level of significance at 0.05

3. RESULTS

The result obtained showed a significantly higher mean value of Plasma Total Cholesterol, LDL cholesterol, Total Triglycerides and HDL cholesterol in rabbits fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days (Group B1 and C1) compared with rabbits fed with normal meal and water for 7 days (Group A) with $p < 0.05$ (Tables 1 and 2, Fig. 1).

There was also a significantly lower mean plasma value of Total Cholesterol, LDL

cholesterol, and Total Triglycerides in rabbits rabbits (B2 and C2) given 400 mg/KgBW of ethanolic and aqueous extract of *Cnidoscopus aconitifolius* for another seven days. Compared with when the same set of rabbits (Group B1 and C1) were only fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days with $p < 0.05$ (Tables 1 and 2, Fig. 1).

There was no significant difference in the Plasma HDL in rabbits fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days (Group B1 and C1) compared to when the same set of rabbits (B2 and C2) were administered with 400 mg/KgBW of ethanolic and aqueous extract of *Cnidoscopus aconitifolius* for another seven days with $p > 0.05$ (Tables 1 and 2, Fig. 1).

Table 1. Mean and standard deviation of the plasma value of total cholesterol, LDL cholesterol, total triglycerides and HDL cholesterol in control and experimental rabbits

	Group A	Group B1	Group B2	Group C1	Group C2
Total cholesterol(mg/dl)	79.0±4.0	122.0±3.0	89.0±1.0	120.0±1.0	90.0±1.0
LDL cholesterol(mg/dl)	42.0±1.0	53.0±3.0	32.0±1.0	51.0±2.0	34.0±1.0
Total triglycerides(mg/dl)	52.0±1.0	76.0±2.0	63.5±0.5	79.0±1.0	63.0±1
HDL cholesterol(mg/dl)	40.0±2.0	54.0 ±3.0	53.0±1.0	59.0±2.0	57.0±1.0

Table 2. Comparative analysis of the plasma value of total cholesterol, LDL cholesterol, total triglycerides and HDL cholesterol in control and experimental rabbits

		A vsB ₁	AVsC ₁	B ₁ Vs, B ₂	C ₁ Vs C ₂	B ₂ Vs C ₂
Total cholesterol(mg/dl)	"t"	-8.6	-9.94	10.44	21.21	-0.71
	"p"	0.007**	0.005**	0.005**	0.001**	0.28
LDL cholesterol(mg/dl)	"t"	-3.48	-3.55	6.64	6.13	-1.41
	"p"	0.04*	0.04*	0.01*	0.02*	0.15
Total Triglycerides(mg/dl)	"t"	-10.7	-19.09	5.81	11.31	0.00
	"p"	0.004**	0.001**	0.01*	0.004**	0.5
HDL cholesterol(mg/dl)	"t"	-3.88	-6.72	0.31	0.89	2.82
	"p"	0.03*	0.01*	0.39	0.23	0.05

* = <0.05-0.01
**= <0.01




Fig. 1. Descriptive analysis of the plasma value of total cholesterol, LDL cholesterol, total triglycerides and HDL cholesterol in control and experimental rabbits

There was no significance difference in plasma value of Total Cholesterol, LDL cholesterol, Total Triglycerides and HDL cholesterol in rabbits hypercholesterolemic rabbits given 400 mg/Kg of aqueous extract of *Cnidoscopus aconitifolius* for another seven days compared with those administered of the ethanolic extract of the same plant with $p > 0.05$ (Tables 1 and 2, Fig. 1).

4. DISCUSSION

The result obtained showed a significantly higher mean value of Plasma Total Cholesterol, LDL cholesterol, Total Triglycerides and HDL cholesterol in rabbits fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days (Group B1 and C1) compared with rabbits fed with normal meal and water for 7 days (Group A).

Significant increase in plasma value of Total Cholesterol, LDL cholesterol, Total Triglycerides and HDL cholesterol following the administration of 20% of powdered egg yolk of the total meal weight in the experimental rabbits could be attributed to the fact that Egg yolk is highly rich in vitamins, minerals, lipids and proteins. It contains nutrient and its primary function is to supply food for the development of the embryo. Egg yolk constitute about 33% of the liquid weight of the egg; it contains about 60 Calories, three times the energy content of the egg white. Furthermore, the weight of egg yolk in a large egg of 50 g total weight, is 17 g which contains about 2.7 g protein, 210 mg cholesterol, 0.61 g carbohydrates, and 4.51 g total fat. It contains fat-soluble vitamins (A, D, E, and K) are found in the egg yolk. Egg yolk, in addition is one of the few foods naturally containing vitamin D [7-10].

There was also a significantly lower mean plasma value of Total Cholesterol, LDL cholesterol, and Total Triglycerides in rabbits rabbits (B2 and C2) given 400 mg/KgBW of ethanolic and aqueous extract of *Cnidoscopus aconitifolius* for another seven days. compared with when the same set of rabbits (Group B1 and C1) were only fed with normal meal containing 20% of powdered egg yolk of the total meal weight and water for seven days This finding could be attributed to the report of NRC [12] that the potassium content (217.2 mg/100 g) is an important mineral nutrient in the control of hypertension and in the reduction of risks of stroke [12]. Rotimi et al. [13] findings also showed that a high-fibre diet modulates plasma lipid and modifies plasma lipoprotein distribution and composition. Albrink et al. [14] reported that

carbohydrate-induced hyperlipemia does not occur if the high carbohydrate diet is rich in dietary fiber, and furthermore that the insulin-stimulating potential of foods in a very high-carbohydrate diet is a critical determinant of the magnitude of carbohydrate-induced lipemia. This could also be attributed to the constituent organic acid (toxic hydrocyanic acid) in the leaf extract [1].

5. CONCLUSION

Induction of hypercholesterolemia using 20% egg yolk of normal meal and water resulted into a significant increase in plasma value of CHOL-T, LDL-C, TG-T, and HDL-C while supplements of 400 mg/kgBW of ethanolic and aqueous extract of *Cnidoscopus aconitifolius* leaf for 7 days significantly reduced the plasma value of CHOL-T, LDL-C and TG-T in the hypercholesterolemic rabbits.

6. RECOMMENDATION

This report is suggested as basis for human trial with respect to hyperlipidemia of hypercholesterolemia.

CONSENT

It is not applicable.

ETHICAL APPROVAL

As per international standard or university standard, written approval of Ethics committee has been collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kuti JO, Konuru HB. Antioxidant capacity and phenolic content in leaf extracts of tree Spinach (*Cnidoscopus* spp.). Journal of Agricultural and Food Chemistry. 2004;52(1):117-21. PMID: 14709023 DOI: 10.1021/jf030246y
2. Ross-Ibarra J, Molina-Cruz A. The ethnobotany of Chaya (*Cnidoscopus aconitifolius* ssp. *Aconitifolius* Breckon): A

- nutritious Maya vegetable. *Economic Botany*. 2002;56(4):350–365. DOI:10.1663/0013-0001(2002)056[0350:TEOCCA]2.0.CO;2 (Retrieved 2007-09-19)
3. Grubben GJH, Denton OA, eds. *plant resources of tropical Africa. 2: Vegetables*. PROTA Foundation. 2004;200–201. ISBN: 978-90-5782-147-9
 4. Kuti JO, Torres ES. Potential nutritional and health benefits of tree spinach. In: J. Janick (ed.), *Progress in New Crops*. ASHS Press, Arlington, VA. 1996;516-520.
 5. Omotoso Abayomi E, Eseyin O Olorunfemi, Suleiman Mikailu. Phytochemical analysis of *Cnidocolus aconitifolius* (Euphorbiaceae) leaf with spectrometric techniques. *Nigerian Journal of Pharmaceutical and Applied Science Research*. 2014;3(1):38-49.
 6. Obichi EA, Monago CC, Belonwu DC. Effect of *Cnidocolus aconitifolius* (Family Euphorbiaceae) aqueous leaf extract on some antioxidant enzymes and haematological parameters of high fat diet and streptozotocin induced diabetic wistar albino rats. *Journal of Applied Sciences and Environmental Management*. 2015; 19(1):201–209.
 7. Richard Stephen Kent. *The invertebrates: A synthesis*. Wiley-Blackwell. 2001;347. ISBN: 978-0-632-04761-1
 8. Bellairs Ruth, Osmond Mark. *Atlas of chick development* (2 ed.). Academic Press. 2005;1-4. Link
 9. Landecker, Hannah. *Culturing life: How cells became technologies*. Cambridge, MA: Harvard University Press. 2007;49. Link
 10. Callebaut M. Historical evolution of preformistic versus neoformistic (epigenetic) thinking in embryology. *Belgian Journal of Zoology*. 2008;138(1): 20–35.
 11. Das K, Tiwari RKS, Shrivastava DK. Techniques for evaluation of medicinal plant products as antimicrobial agent: Current methods and future trends. *Journal of Medicinal Plants Research*. 2010;4(2):104-111.
 12. National Research Council. *Diet and health*. National Academy Press, Washington, DC; 1989.
 13. Rotimi OA, Olayiwola IO, Ademuyiwa O, Balogun EA. Effects of fibre-enriched diets on tissue lipid profiles of MSG obese rats. *Food and Chemical Toxicology*. 2012;50(11):4062.
 14. Albrink MJ, Newman T, Davidson PC. Effect of high- and low-fiber diets on plasma lipids and insulin. *American Journal of Clinical Nutrition*. 1979;32(7): 1486-91.

© 2017 Olaniyan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/20197>